

Guayusa

Ilex guayusa

Family: Aquifoliaceae

By Josef Brinckmann and Thomas Brendler

INTRODUCTION

Ilex guayusa is one of the 500 to 600 *Ilex* species¹ and belongs to the holly family (Aquifoliaceae) of shrubs and trees, which includes only two genera (*Ilex* and *Nemopanthus*).² While the taxonomic status of *Nemopanthus* species remains accepted by some authors, others now consider *Nemopanthus* species to be members of the genus *Ilex*.³ *Ilex* species are distributed in tropical and subtropical to temperate regions, mainly in tropical Central and South America but also Asia.² The perennial *I. guayusa* tree, native to Bolivia, Colombia, Ecuador, Peru, and Venezuela,⁴ is dioecious (male and female flowers occur on separate plants) and reaches about 10 m (32.8 ft) when cultivated and up to 25 m (82 ft) when unmanaged or wild.⁵ Other authors give a range of 6-30 m (19.7-98.4 ft). Guayusa is fast-growing, and commercial harvesting of the leaves may commence about one year after planting, with increasing annual yields until leveling off with tree maturity at around five years.⁶

Domesticated in the pre-Columbian era, guayusa is now grown in traditional agroforestry systems by indigenous farmers of the western Amazon region. In recent years, scaled-up production of certified organic and fair trade guayusa leaf has linked some indigenous farmers in the Ecuadorian Amazon to a global market for sustainable herbs, helping to conserve biodiversity (prevent forest clearing) and improve traditional economies in rural and remote indigenous communities.⁶ Emerging evidence suggests that traditional agroforestry systems support climate change adaptation because they enable measurably higher levels of carbon sequestration and tree diversity compared to other forms of cultivation.⁷

HISTORY AND CULTURAL SIGNIFICANCE

Ilex guayusa is known as *waysa*⁷ or *guayusa* in Kichwa, a dialect of the Quechua language group, the latter also used by Spanish speakers,⁸ and as *wais* in the Shuar language in Ecuador.⁹ A standard common name for this species is not provided in the American Herbal Products Association's *Herbs of Commerce*, 2nd edition.¹⁰ In his 1737 work *Genera Plantarum*, Swedish botanist Carl Linnaeus (1707-1778) assigned the genus name *Ilex*,¹¹ which was already the Latin name for the Mediterranean holly oak (*Quercus ilex*, Fagaceae), probably due to the similar glossy leaves.¹² German botanist Ludwig Eduard Theodor Loesener (1865-1941) assigned the Kichwa name guayusa as the species name, forming the Latin binomial *Ilex guayusa*, in his 1901 publication *Monographia Aquifoliacearum*.¹³

Archaeobotanical evidence suggests that guayusa has been used medicinally and traded since at least 500 BCE in the greater Andes-Amazon region.⁵ The contents of a tomb dated to that time, found at Niño Korin in the Bautista Saavedra Province of Bolivia, included pouches of guayusa leaves as well as *Anadenanthera colubrina* (Fabaceae) along with specialized medical equipment (e.g., enema syringes and snuff trays) believed to have belonged to a medicine person of the pre-Incan Kallawayaya society of traditional healers, descendants of the Tiwanaku culture.¹⁴ Although cultivation may have begun earlier, evidence suggests that guayusa was grown extensively and processed by at least 350 CE.¹⁵ In 1972, American renowned Harvard University ethnobotanist Richard Evans Schultes, PhD, (1915-2001) reported discovering the vestiges of an ancient guayusa plantation in Putumayo Department of southwestern Colombia, with tall trees that were hundreds of years old.¹⁶

A 1683 letter from Jesuit priest Juan Lorenzo Lucero to the Viceroy of Peru, Don Melchor de Navarra y Rocafull,



Guayusa *Ilex guayusa*

Photo ©2019 Applied Food Sciences

described the drinking of a decoction of guayusa throughout the day by the indigenous Jivaroan peoples of Ecuador and Peru for the purpose of remaining awake for many nights at times when an enemy invasion was feared.¹⁷ Seventeenth-century Jesuit missionaries planted guayusa in mission gardens and established commercial trade to highland Andean markets, including Quito.¹⁸ According to Schultes, the Jesuits also established a market for guayusa leaf in Europe as a cure for syphilis, among other diseases.¹⁹ With the expulsion of the Jesuits from the Spanish colonies in the 1760s, small-scale cultivation and regional trade were taken over by indigenous peoples, and guayusa is still found today in Ecuadorian village markets.¹⁸ In 1894, anthropologist Charles Dolby Tyler wrote that guayusa, also known as “Napo tea,” was the favorite beverage of the Záparo people, who lived along the Napo River in Ecuador. The Záparo drank the tea in the morning as an emetic to rid the stomach of undigested food. Tyler also noted their use of coca (*Erythroxylum coca*, Erythroxylaceae) leaf, which was chewed and mainly taken before long journeys.²⁰

Shamans of the indigenous Shuar people in Amazonian Ecuador and Peru include guayusa in hallucinogenic herbal formulations, although it is a stimulant rather than a hallucinogen. For example, guayusa is combined with *Banisteriopsis caapi* (known by the local common name *ayahuasca*)

and *Diplopterys cabrerana* (both in the family Malpighiaceae) and prepared as an aqueous decoction.⁹ Guayusa tea is also reportedly used by Canelos-Kichwa people before and after drinking ayahuasca tea (i.e., the blend of *Banisteriopsis caapi* and other plants), as it is believed to mitigate the bitter taste of the ayahuasca, prevent hangover, and strengthen the ability to cope with the hallucinogenic effects.²¹ As a daily ritual, men of the Jivaroan Achuar community in Ecuador and Peru drink aqueous decoctions of guayusa leaf, called *wayus*, before sunrise, and then vomit, for stimulating effects. Boys are taught how to vomit using a feather or finger, and they join the men in the daily ritual after puberty.^{22,23*} A study looking at obstetrical practices of Canelos-Kichwa women living in communities along the Bobonaza River in the Pastaza Province of Ecuador, near the Peruvian border, found that guayusa tea is drunk for the purpose of reducing blood loss during menstruation, with concomitant avoidance of both salt and chili (*Capsicum* spp., Solanaceae) for one to three days. Postpartum, women are also given guayusa tea to drink in order to “prevent blood drying up inside her and thus cause other complications.”²⁴

In a study of traditional plant usage among the indigenous Kichwa people of Canton Loreto in Ecuador, guayusa was found to be the most used, harvested, and significant plant in daily life.²⁵ In another recent study that surveyed indigenous Shuar and *mestizos* about traditional ecological knowledge and medicinal plant diversity in Ecuadorian Amazon home gardens, guayusa ranked among the highest in importance. The study asserts that guayusa and other medicinal plants grown in these home gardens are ecosystem service providers that support human well-being, biodiversity conservation, and traditional knowledge in agroecosystems.²⁶

CURRENT AUTHORIZED USES IN COSMETICS, FOODS, AND MEDICINES

In the United States, guayusa is marketed as a beverage tea for energy, similar to caffeine-containing leaves of other *Ilex* species such as yerba maté (*I. paraguariensis*) and yaupon (*I. vomitoria*).²⁷ It may also be used as a component of dietary supplement products, which require notification with the US Food and Drug Administration within 30 days of marketing if a structure-function claim is made.²⁸ In Canada, guayusa is regulated as a medicinal ingredient of licensed natural health products (NHPs), which require pre-marketing authorization from the Natural and Non-prescription Health Products Directorate (NNHPD).²⁹ At the time of this writing, two licensed NHPs in Canada list *I. guayusa* as an active ingredient.³⁰ In the EU, use of

* Although English botanist William Townsend Aiton (1766–1849) assigned the Latin binomial *Ilex vomitoria* to the related North American yaupon,⁶⁶ and both guayusa and yaupon have been used traditionally in indigenous vomiting rituals,⁶⁷ constituents that could be responsible for emetic action have not been identified in either species.⁶⁸





Guayusa *Ilex guayusa*
Photo ©2019 Applied Food Sciences

an extract of the leaves in cosmetic products is authorized specifically for skin-protecting function.³¹ However, for oral ingestion, *I. guayusa* is presently classified as an unauthorized novel food in the EU. According to EU regulatory authorities, *I. guayusa* was not used as a food or food ingredient in the EU before May 15, 1997, and, therefore, a pre-marketing safety assessment under the Novel Food Regulation is required.³²

MODERN RESEARCH

Largely seen as a stimulant traditional beverage among indigenous communities,^{9,16,17,22} with some medicinal uses therein (e.g., to avoid postpartum complications; as a tonic, diuretic, flu remedy, mouth wash, and insect repellent; for dysmenorrhea, diabetes, venereal diseases, and weight loss^{5,24,26,33-35}), guayusa only recently attracted interest from the research community for its health benefits. Published research on its actions and effects remains limited. Many recent academic theses investigating medicinal and cosmetic applications of guayusa³⁶⁻⁴² indicate new interest, likely also stimulated by international commercialization and regulatory requirements.^{35,43,44} There are no known human clinical trial publications to date.

Plants in the genus *Ilex* are known to contain a significant number of secondary metabolites, such as xanthines, chlorogenic acid derivatives, flavonoids, and triterpenoids.^{33,34,45,46} Wise and Santander (2018) provide a comprehensive summary of the composition of dried guayusa leaves, including total and free amino acids, chemical

elements, nutritional values, and caffeine content.⁴⁷ Earlier research addressing the composition of guayusa focused on its high caffeine content (e.g., Lewis et al., 1991).²² More recently, triterpenoids in guayusa leaf, specifically ursolic acid, have gained attention due to their antihyperglycemic, antibacterial, and antiparasitic effects.³³⁻³⁵

Ursolic acid is a known activator of the G protein-coupled bile acid receptor 1 (GPBAR1), which is involved in energy homeostasis, bile acid homeostasis, glucose metabolism, inflammatory response, cancer progression, and liver regeneration, giving guayusa a potential role in the management of diabetes and metabolic syndrome. Swanston-Flatt et al. (1989, 1991)^{48,49} showed that guayusa had a hypoglycemic effect in normal mice and in mice with streptozotocin-induced diabetes. In addition, guayusa reduced hyperphagia (increased appetite), polydipsia (excessive thirst), body weight loss, and glycated hemoglobin. These results were confirmed by Stoyell-Conti et al. (2018),⁵⁰ who showed positive effects on metabolic, cardiovascular, and oxidative stress in mice with streptozotocin-induced diabetes. Espinosa Soto et al. (2015)⁵¹ reported the presence of ursolic acid, which was confirmed by Chianese et al. (2019),⁵² who determined a similar triterpenoid profile for *I. guayusa* and *I. paraguayensis*, as well as the presence of amyirin esters. They characterized amyirin palmitate, palmitoleate, and corresponding isomers as the primary constituents of the amyirin complex from both plants.

García-Ruiz et al. (2017)⁵³ identified a total of 14 phenolic compounds and five carotenoids in guayusa leaf, and

further showed high antioxidant capacity for blanched and untreated leaves. This antioxidant capacity was much reduced during fermentation. Pardau et al. (2017)⁵⁴ demonstrated protection from oxidative stress in the Caco-2 cellular antioxidant assay and anti-inflammatory activity in lipopolysaccharide-stimulated RAW 264.7 cells, and these effects were attributed to the phenolic mono- and dicaffeoylquinic acid derivatives in guayusa leaves. Cadena-Carrera et al. (2019)⁵⁵ studied the biological activities of guayusa leaf extracts using supercritical CO₂ as the solvent and ethanol as the co-solvent and showed antifungal activity against *Trichophyton rubrum*, *T. mentagrophytes*, *Microsporum gypseum*, and *M. canis*. Caffeine, squalene, and α -amyirin were the main compounds found; antioxidant activity varied with extraction technique, solvent, and conditions (e.g., pressure, temperature). Gamboa et al. (2018)⁵⁶ demonstrated an antimicrobial effect for an ethanolic extract of guayusa leaves, inhibiting *Porphyromonas gingivalis*, *Prevotella intermedia*, and *Fusobacterium nucleatum*, bacteria implicated in chronic periodontitis. Both Tuquinga Usca (2013)⁵⁷ and Contero et al. (2015)⁵⁸ reported an estrogenic effect of guayusa in albino rats, both in terms of significantly increased estradiol levels and weight of reproductive organs.

Sequeda-Castañeda et al. (2016)³³ summarized a number of investigations of the safety of guayusa preparations and attest an excellent safety profile. No signs of hepatotoxicity were found in an in vivo hepatotoxicity model using Wistar rats. No acute toxicity was shown at 1,000, 500, 250, and 125 mg/kg ethanol extract in animals. Repeat doses were also found to be safe. Due to the high caffeine content, however, large quantities can affect the nervous system. Bussmann et al. (2011)⁵⁹ demonstrated a LC₅₀ > 10,000 μ g/mL for an aqueous extract and 300 μ g/mL for an ethanol extract of guayusa leaves in the brine shrimp lethality assay. Kapp et al. (2016)⁶⁰ found guayusa concentrate to be negative in in vitro genotoxicity tests, including the Ames test and a chromosome aberration study in human lymphocytes. This confirmed previous results finding an LD₅₀ > 5,000 mg/kg for female rats. A 90-day sub-chronic study at 1,200, 2,500, and 5,000 mg/kg/d of guayusa concentrate administered to male and female rats found effects comparable to those of caffeine, including weight loss, reductions in food efficiency and triglycerides values, increases in serum alanine aminotransferase, serum aspartate aminotransferase, and cholesterol, as well as adaptive salivary gland hypertrophy. Overall, no harmful effects specific to guayusa or its components were observed in any of the tested models.



Tea being made from guayusa (*Ilex guayusa*)
Photo ©2019 Applied Food Sciences

ADULTERATION

Although adulteration with other related plants, such as yerba maté, is conceivable, adulteration of guayusa products is, so far, not known to occur. While there is a *European Pharmacopoeia* monograph for testing of yerba maté (Mate Folium PhEur 10.0),⁶¹ which provides macroscopic, microscopic, thin-layer-chromatography (TLC), and high-performance-liquid-chromatography (HPLC) tests for confirming composition, identity, quality, and strength, no known monographs are available for guayusa or yaupon that provide identification methods for differentiating or ruling out admixing among the three related species.

SUSTAINABILITY AND FUTURE OUTLOOK

The International Union for Conservation of Nature (IUCN) assigns wild *I. guayusa* to the conservation category of least concern (LC), meaning that the species is not considered to be threatened. According to the 2019 IUCN report, “this species has a very wide distribution, large population, is not currently experiencing any major threats, and no significant future threats have been identified.”⁴

In 2009, the company RUNA (Brooklyn, New York) began to commercialize guayusa in Ecuador and, by 2017, had 477 hectares of cultivation area under US Department

of Agriculture (USDA) National Organic Program (NOP) certification.⁶ Today, NOP-certified guayusa operations in Ecuador include Asociación de Caficultores el Pangui, Fundación Chankuap “Recursos Para el Futuro,” Greenmattersecuador CIA LTDA, Jumandipro S.A., Productos SKS Farms CIA LTDA, Rarecats S.A., Runatarpuna Exportadora S.A., and Tryskelwork Soluciones S.A.⁶² One of these organic guayusa operations has also achieved economic and social sustainability certification through implementation of the Fair Trade USA standards: Runatarpuna Exportadora S.A. (Quito, Pichincha, Ecuador), which exports organic- and fair trade-certified guayusa to their North American counterpart, RUNA. Runatarpuna’s guayusa farmers are situated primarily in Napo Province in the Ecuadorian Amazonian rainforest, and cultivate it within a multi-crop traditional agroforestry system along with cacao (*Theobroma cacao*, Malvaceae), coffee (*Coffea arabica*, Rubiaceae), and yuca (*Manihot esculenta*, Euphorbiaceae).⁶³ As opposed to mono-cropping, these traditional agroforestry practices are believed to mitigate some of the local effects of climate change and increase food security for smallholder farmers.⁶

RUNA is also a Certified B Corporation, audited under a certification standard that measures a company’s social and environmental performance, public transparency, and legal accountability.⁶⁴ In 2018, RUNA’s nonprofit arm, Runa Foundation, merged with PlanJunto Ltda to form a new organization called Aliados, whose mission is “to build resilient community business based on biodiversity in the Andes and the Amazon — and connect them to markets across the globe.” A current Aliados venture involves a partnership with Ally Guayusa, an indigenous-owned guayusa export association composed of 140 farming families operating 40 hectares of organic guayusa production.⁶⁵


Before RUNA was founded in 2009, the use of guayusa leaf was relatively unknown outside of the Amazon. The socially conscious and responsible manner in which RUNA has sought to bring guayusa to the world is admirable. In collaboration with the local farming communities, RUNA has invested in building the foundation for a guayusa market that is ecologically, economically, and socially sustainable. However, Krause and Ness (2017) caution that guayusa production is still a niche agroforestry experiment and that the sustainability initiatives, such as fair trade and organic standards and certifications, so far “only provide partial solutions for protecting ecosystem services in the Ecuadorian Amazon.” Growing popularity and success of guayusa in the global market may present the risk of its becoming a new monoculture cash crop that could displace diverse agroforestry systems.⁶ HG

Declaration of Interest

The authors declare that they have no conflict of interest and no affiliation with the Runa organizations.



30 years
of expertise
in phytochemical
reference
substances



Cranberry – Procyanidin A2

www.extrasynthese.com

True standards for identification, adulteration tracking
and quantification of botanicals and herbal products.



**NOT
GINSENG**

Root, Leaf, or Flower?

Plant part adulteration is an unfortunate reality. Are you getting what you paid for? Is it really what you are claiming on your label? We're here to help. No other contract lab knows herbal testing like Alkermist.



ALKEMIST LABS

714.754.4372 ALKEMIST.COM

References

- Cuénoud P, Del Pero Martínez MA, Loizeau P-A, Spichiger R, Andrews S, Manen J-F. Molecular phylogeny and biogeography of the genus *Ilex* L. (Aquifoliaceae). *Ann Bot*. 2000;85(1):111-122.
- Chen S, Ma H, Feng Y, Barrera G, Loizeau PA. Aquifoliaceae. 冬青科 dong qing ke. In: *Flora of China*. Vol 11. St. Louis, MO: Missouri Botanical Garden Press; 2008:359-438.
- NatureServe. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. Arlington, VA: NatureServe; 2019.
- Botanic Gardens Conservation International (BGCI) & IUCN SSC Global Tree Specialist Group. *Ilex guayusa*. The IUCN Red List of Threatened Species 2019: eT145566259A145678088 <http://dxdoiorg/102305/IUCN2019-2RLTS-T145566259A145678088en>. 2019.
- Dueñas JF, Jarrett C, Cummins I, Logan-Hines E. Amazonian guayusa (*Ilex guayusa* Loes.): A historical and ethnobotanical overview. *Econ Bot*. 2016;70(1):85-91.
- Krause T, Ness B. Energizing agroforestry: *Ilex guayusa* as an additional commodity to diversify Amazonian agroforestry systems. *Int J Biodivers Sci Ecosyst Serv Manag*. 2017;13(1):191-203.
- Torres B, Maza OJ, Aguirre P, Hinojosa L, Günter S. The Contribution of Traditional Agroforestry to Climate Change Adaptation in the Ecuadorian Amazon: The Chakra System. In: Leal Filho W, ed. *Handbook of Climate Change Adaptation*. Berlin, Heidelberg, Germany: Springer Berlin Heidelberg; 2015:1973-1994.
- Edwards AL, Bennett BC. Diversity of methylxanthine content in *Ilex cassine* L. and *Ilex vomitoria* Ait.: Assessing sources of the North American stimulant cassina. *Econ Bot*. 2005;59(3):275-285.
- Bennett BC. Hallucinogenic plants of the Shuar and related indigenous groups in Amazonian Ecuador and Peru. *Brittonia*. 1992;44(4):483-493.
- McGuffin M, Kartesz JT, Leung AY, Tucker AO. *American Herbal Products Association's Herbs of Commerce*. 2nd ed. Silver Spring, MD: American Herbal Products Association; 2000.
- Linné Cx. *Caroli Linnæi ... Genera plantarum eorumque characteres naturales secundum numerum, figuram, situm, & proportionem omnium fructificationis partium*. Lugduni Batavorum :: apud C. Wishoff; 1737.
- Marzell H. *Wörterbuch der deutschen Pflanzennamen. 2. Band*. Stuttgart/Wiesbaden: S. Hirzel Verlag/Franz Steiner Verlag; 1977.
- Loesener T. Monographia Aquifoliacearum. Pt. I. In: *Abhandlungen der Kaiserlichen Leopoldinisch-Carolinischen Deutschen Akademie der Naturforscher*. Vol 78. Leipzig, Preußen: W. Engelmann; 1901:311.
- Wassén H. *A Medicine-Man's Implements and Plants in a Tiahuanaoid Tomb in Highland Bolivia. Etomologica studier*, Vol. 32. Göteborg, Sweden: Göteborgs Etnografiska Museum; 1972.
- Lathrap DW. The antiquity and importance of long-distance trade relationships in the moist tropics of pre-Columbian South America. *World Archaeology*. 1973;5(2):170-186.
- Schultes RE. Discovery of an ancient guayusa plantation in Colombia. *Botanical Museum Leaflets, Harvard University*. 1979;27(5/6):143-153.
- Patiño VM. Guayusa, a neglected stimulant from the eastern Andean foothills. *Econ Bot*. 1968;22(4):310-316.
- Jamieson RW. The essence of commodification: Caffeine dependencies in the Early Modern World. *J Soc Hist*. 2001;35(2):269-294.
- Schultes RE. Antiquity of the use of new world hallucinogens. *The Heffter Review of Psychedelic Research*. 1998;1:1-7.
- Tyler CD. The River Napo. *The Geographical Journal*. 1894;3(6):476-484.
- Shemluck M. The flowers of *Ilex guayusa*. *Botanical Museum Leaflets, Harvard University*. 1979;27(5/6):155-160.
- Lewis WH, Kennelly EJ, Bass GN, Wedner HJ, Elvin-Lewis MP, W DF. Ritualistic use of the holly *Ilex guayusa* by Amazonian Jivaro Indians. *J Ethnopharmacol*. 1991;33(1):25-30.
- Lewis WH, Elvin-Lewis MP. Medicinal plants as sources of new therapeutics. *Ann Miss Bot Gard*. 1995;82(1):16-24.
- Baruffati V. The reproductive cycle of Ecuadorian rain forest women. *Bull Lat Am Res*. 1984;3(1):99-109.
- Innerhofer S, Bernhardt K-G. Ethnobotanic garden design in the Ecuadorian Amazon. *Biodivers Conserv*. 2011;20(2):429-439.
- Caballero-Serrano V, McLaren B, Carrasco JC, et al. Traditional ecological knowledge and medicinal plant diversity in Ecuadorian Amazon home gardens. *Glob Ecol Cons*. 2019;17:e00524.
- Runa. Clean energy® drink. Available at: <https://runa.com/our-story>. Accessed August 11, 2019.
- US Food and Drug Administration. Regulations on Statements Made for Dietary Supplements Concerning the Effect of the Product on the Structure or Function of the Body; Final Rule. *Federal Register*. 2000;65(4):1000-1050.
- Natural and Non-prescription Health Products Directorate. *Natural Health Products Ingredients Database*. Ottawa, Ontario: Health Canada; 2019.
- Natural and Non-prescription Health Products Directorate. *Licensed Natural Health Products Database (LNHPD)*. Ottawa, Ontario: Health Canada; 2019.
- European Commission. *Cosmetic ingredient (CosIng) database*. Brussels, Belgium: DG Internal Market, Industry, Entrepreneurship and SMEs; 2019.
- European Commission. *Novel Food Catalogue*. Brussels, Belgium: European Commission; 2019.
- Sequeda-Castañeda L, Costa G, Celis C, Gamboa F, Gutierrez S, Luengas P. *Ilex guayusa* (Aquifoliaceae): Amazon and Andean native plant. *Pharmacologyonline*. 2016;3:93-202.
- Radice M, Scalvenzi L, Sablón Cossío N. *Ilex guayusa*: A systematic review of its traditional uses, chemical constituents, biological activities and biotrade opportunities. In: *Proceedings of MOL2NET 2016, International Conference on Multidisciplinary Sciences*. 2nd ed. Basel, Switzerland: MDPI; 2017:3868.
- Wise G, Negrin A. A critical review of the composition and history of safe use of guayusa: a stimulant and antioxidant novel food. *Crit Rev Food Sci Nutr*. 2019:1-12.
- Calderón Borja EA. *Investigación y desarrollo de un protector solar empleando como ingrediente activo los agentes antioxidantes en extractos de guayusa (Ilex guayusa Loes)*. Quito, Ecuador: Universidad San Francisco de Quito, Colegio de Ciencias e Ingenierías; 2017.
- Espinosa Caiza EP. *Actividad anti-biofilm de los extractos de plantas Urtica dioica L., Ilex guayusa y Uncaria tomentosa en Staphylococcus aureus*. Universidad Técnica de Ambato - Facultad de Ciencias de la Salud - Carrera de Laboratorio Clínico. 2019.
- Carpintero Salvador N, Salazar Enríquez ME. *Evaluación del efecto anticelulítico de una formulación cosmética a base de extracto alcohólico foliar de Guayusa, Ilex guayusa Loes (Aquifoliaceae)*. Quito, Ecuador: Universidad Politécnica Salesiana; 2014.
- Hernández Dávila CA. *Elaboración y caracterización de Microcápsulas mediante Gelificación Iónica externa, de la fracción alcaloidea de ILEX GUAYUSA con ALGINATO SÓDICO, y su utilización en la formulación de una forma de dosificación sólida*. Escuela Superior Politécnica de Chimborazo; 2015.
- Silva Escobar ME. *Valoración de las especies Ilex guayusa (GUAYUSA) y Plukenetia volubilis L. (SACHA INCHI), como una materia prima cosmética antioxidante*. Quito, Ecuador: Universidad Politécnica Salesiana; 2018.
- Pardau MD. *Antioxidant content activity and nitric oxide scavenging ability of Ilex guayusa compared with Camellia sinensis and Aspalathus linearis tea*. MSc Dissertation, University of Pretoria, Pretoria, South Africa; 2016.
- Palchizaca Pinguil MN. *Evaluación del efecto antiinflamatorio de Ilex guayusa (Loes), Vernonia patens (Kuntz) y Theobroma cacao (Linneo) en el modelo Animal Rata (Rattus norvegicus) de laboratorio*. Escuela Superior Politécnica de Chimborazo. Riobamba; 2019.
- Wise G. *Substantial Equivalence Application for the Approval of Guayusa Leaf Tea (Ilex guayusa Loes.) for Use as a Dry Leaf Infusion*. Tena, Ecuador: IKIAM EP; 2017. Available at: <https://acnfp.food.gov.uk/sites/default/files/guayusaapp.pdf>. Accessed September 9, 2019.

44. Runa LCC. Substantial Equivalence of Aqueous Extracts of the Dried Leaves of *Ilex guayusa*. As cited in Food Safety Authority of Ireland Substantial Equivalence Opinion *Ilex guayusa*. 2017. Available at: www.fsai.ie/uploadedFiles/Science_and_Health/Novel_Foods/Notifications/2017%20SE%20Opinion%20Ilex%20Guayusa.pdf. Accessed September 9, 2019.
45. Yi F, Zhao X-l, Peng Y, Xiao P-g. Genus *Ilex* L.: Phytochemistry, ethnopharmacology, and pharmacology. *Chin Herb Med*. 2016;8(3):209-230.
46. Gan R-Y, Zhang D, Wang M, Corke H. Health benefits of bioactive compounds from the genus *Ilex*, a source of traditional caffeinated beverages. *Nutrients*. 2018;10(11):1682.
47. Wise G, Santander DE. Comparative composition analysis of the dried leaves of *Ilex guayusa* (Loes.). *J Food Nutr Res*. 2018;6(10):638-644.
48. Swanston-Flatt SK, Day C, Flatt PR, Gould BJ, Bailey CJ. Glycaemic effects of traditional European plant treatments for diabetes. Studies in normal and streptozotocin diabetic mice. *Diabetes Res*. 1989;10(2):69-73.
49. Swanston-Flatt SK, Flatt PR, Day C, Bailey CJ. Traditional dietary adjuncts for the treatment of diabetes mellitus. *Proc Nutr Soc*. 1991;50(3):641-651.
50. Stoyell-Conti FF, Mello TC, Dias DS, et al. Effects of *Ilex guayusa* tea on glycaemia, cardiovascular autonomic modulation and oxidative stress in streptozotocin-diabetic rats. *The FASEB Journal*. 2018;32(1_supplement):906-912.
51. Espinosa Soto I, Valle ME, Menestrina B, Rosella MA. PS029: Ursolic acid in *Ilex guayusa* Loes. (Aquifoliaceae). In: *XXIV SILAE Congress. II International Congress on Integrative Medicine. Congress Abstract Book*. Salerno, Italy: Società Italo-Latino Americana di Etnomedicina (SILAE); 2015:97.
52. Chianese G, Golin-Pacheco SD, Tagliatalata-Scafati O, et al. Bioactive triterpenoids from the caffeine-rich plants guayusa and maté. *Food Res Int*. 2019;115:504-510.
53. García-Ruiz A, Baenas N, Benítez-González AM, et al. Guayusa (*Ilex guayusa* L.) new tea: Phenolic and carotenoid composition and antioxidant capacity. *J Sci Food Agric*. 2017;97(12):3929-3936.
54. Pardau MD, Pereira ASP, Apostolides Z, Serem JC, Bester MJ. Antioxidant and anti-inflammatory properties of *Ilex guayusa* tea preparations: A comparison to *Camellia sinensis* teas. *Food Funct*. 2017;8(12):4601-4610.
55. Cadena-Carrera S, Tramontin DP, Bella Cruz A, Bella Cruz RC, Müller JM, Hense H. Biological activity of extracts from guayusa leaves (*Ilex guayusa* Loes.) obtained by supercritical CO₂ and ethanol as cosolvent. *J Supercrit Fluid*. 2019;152:104543.
56. Gamboa F, Muñoz C-C, Numpaque G, Sequeda-Castañeda LG, Gutierrez SJ, Tellez N. Antimicrobial activity of *Piper marginatum* Jacq and *Ilex guayusa* Loes on microorganisms associated with periodontal disease. *Int J Microbiol*. 2018;2018(Article ID 4147383):9.
57. Tuquinga Usca MR. *Efecto Estrogénico del Extracto de las Hojas de Guayusa (Ilex guayusa Loes) en Ratas (Rattus norvegicus)*. Escuela Superior Politécnica de Chimborazo; 2013.
58. Contero F, Abdo S, Vinuza D, Moreno J, Tuquinga M, Paca N. Estrogenic activity of ethanolic extracts from leaves of *Ilex guayusa* Loes. and *Medicago sativa* in *Rattus norvegicus*. *Pharmacologyonline*. 2015;2:95-99.
59. Bussmann RW, Malca G, Glenn A, et al. Toxicity of medicinal plants used in traditional medicine in Northern Peru. *J Ethnopharmacol*. 2011;137(1):121-140.
60. Kapp RW, Mendes O, Roy S, McQuate RS, Kraska R. General and genetic toxicology of guayusa concentrate (*Ilex guayusa*). *Int J Toxicol*. 2016;35(2):222-242.
61. European Pharmacopoeia Commission. *European Pharmacopoeia*. 10th ed. (PhEur 10.0). Strasbourg, France: European Directorate for the Quality of Medicines; 2019.
62. United States Department of Agriculture. *Organic INTEGRITY Database*. Washington, D.C.: USDA Agricultural Marketing Service; 2019.
63. Nealis K. Trip report: Fair trade impact in Ecuador. Fair Trade Campaigns Blog. 2017. Available at: <https://fairtradecampaigns.org/2017/08/trip-report-fair-trade-impact-in-ecuador/>. Accessed August 11, 2019.
64. B Lab. Certified B Corporation. B Impact Report: Runa. Available at: <https://bcorporation.net/directory/runa>. Accessed August 17, 2019.
65. Aliados. Ally Guayusa: Organic guayusa tea. Available at: www.losalidos.org/ally-guayusa. Accessed August 17, 2019.
66. Alston AHG, Schultes RE. Studies of early specimens and reports of *Ilex vomitoria*. *Rhodora*. 1951;53(636):273-279.
67. Samorini G. The oldest archeological data evidencing the relationship of *Homo sapiens* with psychoactive plants: A worldwide overview. *J Psyched Stud*. 2019;3:1-18.
68. McClatchey WC, Mahady GB, Bennett BC, Shiels L, Savo V. Ethnobotany as a pharmacological research tool and recent developments in CNS-active natural products from ethnobotanical sources. *Pharmacol Ther*. 2009;123(2):239-254.

ABC Board of Trustees

Michael J. Balick, PhD

VP and Director,
Institute of Economic Botany, New York
Botanical Garden, Bronx, NY

Neil Blomquist

President, Sustainable Solutions
Consulting Services, Sebastopol, CA

Peggy Brevoort

President, Brevoort, LLC, Kapa'au, HI

Bethany Davis

Director, Regulatory & Industry Affairs
FoodState Inc., Manchester, NH

Steven Foster

President, Steven Foster Group, Inc.,
Eureka Springs, AR

Richard Kingston, PharmD, CSPI

President, Regulatory and Scientific Affairs,
Safety Call International Poison Center;
Professor, Dept. of Experimental & Clinical
Pharmacology, University of Minnesota,
Minneapolis, MN

Bernadette P. Marriott, PhD

Professor, Department of Medicine
MUSC College of Medicine
Div. of Gastroenterology & Hepatology &
Department of Psychiatry
and Behavioral Sciences
Military Division, Charleston, SC

Thomas M. Newmark

Owner, Finca Luna Nueva Lodge, San
Isidro de Peñas Blancas, Costa Rica

Holly Shimizu

Consultant, Writer, and Lecturer
Former Executive Director, US Botanic
Garden; First Curator, The National Herb
Garden, Glen Echo, MD

Morris Shriftman

CEO, Mozart, Inc., Petaluma, CA

Margaret Wittenberg

Retired, Natural & Organic Foods Industry
Fredericksburg, TX

Mark Blumenthal

(*ex officio*)
Founder &
Executive Director
American
Botanical Council
Austin, TX

